arXiv:1303.5527v1 [math.CO] 22 Mar 2013

# Signless laplacian characteristic polynomials of regular graph transformations

Jianping Li[a,b], Bo Zhou[b*]
[a]Faculty of Applied Mathematics, Guangdong University of Technology,
Guangzhou 510090, P. R. China
[b]Department of Mathematics, South China Normal University,
Guangzhou 510631, P. R. China
Email: `zhoubo@scnu.edu.cn`

**Abstract**

Let $G$ be a simple $r$-regular graph with $n$ vertices and $m$ vertices. We give the signless Laplacian characteristic polynomials of $xyz$-transformations $G^{xyz}$ of $G$ in terms of $n$, $m$, $r$ and the signless Laplacian spectrum of $G$.

**Keywords:** regular graph, $xyz$-transformation, signless Laplacian characteristic polynomial

## 1 Introduction

We consider simple graphs. Let $G = (V, E)$ be a graph with vertex set $V(G)$ and edge set $E(G)$.

Let $V(G) = \{v_1, \ldots, v_n\}$. The adjacency matrix $A(G)$ of $G$ is the $(0,1)$-matrix $(a_{ij})$ of order $n$ where $a_{ij} = 1$ if $v_iv_j \in E(G)$, and $a_{ij} = 0$ otherwise. The degree matrix $D(G)$ of $G$ is the (diagonal) matrix $(d_{ij})$ of order $n$ where $d_{ii}$ is the degree of vertex $v_i$ in $G$ and $d_{ij} = 0$ for $i \neq j$. The matrix $L(G) = D(G) - L(G)$ is the Laplacian matrix of $G$, and $Q(G) = D(G) + A(G)$ is the signless Laplacian matrix of $G$.

---

[*]Corresponding author.

The characteristic polynomials of $A(G)$, $L(G)$ and $Q(G)$ are called the adjacency, Laplacian and signless Laplacian characteristic polynomials of $G$, respectively. The spectra of $A(G)$, $L(G)$ and $Q(G)$ are called the adjacency, Laplacian and signless Laplacian spectra of $G$, respectively.

The complement $G^c$ of $G$ is the graph with vertex set $V(G^c) = V(G)$ and for any $u, v \in V(G)$ and $u \neq v$, $uv \in E(G^c)$ if and only if $uv \notin E(G)$ .

Let $G^0$ be the empty graph with $V(G^0) = V(G)$, $G^1$ the complete graph with $V(G^1) = V(G)$, $G^+ = G$, and $G^- = G^c$.

Let $B(G)$ ($B^c(G)$) be the graph with vertex set $V(G) \cup E(G)$ such that $ve$ is an edge in $B(G)$ (resp., in $B^c(G)$) if and only if $v \in V(G)$, $e \in E(G)$, and vertex $v$ is incident (resp., not incident) to edge $e$ in $G$.

The line graph $G^l$ of $G$ is the graph with vertex set $E(G)$ and two vertices are adjacent in $G^l$ if and only if the corresponding edges in $G$ are adjacent.

Let $G$ be a graph and $x, y, z$ variables in $\{0, 1, +, -\}$. The $xyz$-transformation $G^{xyz}$ of $G$ is the graph with vertex set $V(G^{xyz}) = V(G) \cup E(G)$ and the edge set $E(G^{xyz}) = E(G^x) \cup E((G^l)^y) \cup E(W)$, where $W = B(G)$ if $z = +$, $W = B^c(G)$ if $z = -$, $W$ is the empty graph with $V(W) = V(G) \cup E(G)$ if $z = 0$, and $W$ is the complete bipartite graph with partite sets $V(G)$ and $E(G)$ if $z = 1$.

For a regular graph $G$, the adjacency characteristic polynomials (and the adjacency spectra) of $G^{00+}$, $G^{+0+}$, $G^{0++}$ and $G^{+++}$ can be found in [1]. The adjacency characteristic polynomials (and the adjacency spectra) of the other seven $G^{xyz}$ with $x, y, z \in \{+, -\}$ can be found in [3]. Deng et al. [4] determined the Laplacian characteristic polynomials of $G^{xyz}$ of a regular graph $G$ with $x, y, z \in \{0, 1, +, -\}$ (The cases $G^{0++}$, $G^{0+0}$ and $G^{00+}$ have early been given in[2]).

Let $f(x, G) = \det(xI_n - Q(G))$ be the signless Laplacian characteristic polynomial of $G$, where $I_n$ is the the identity matrix of order $n = |V(G)|$. Now we give the signless Laplacian characteristic polynomials of $xyz$-transformation of an $r$-regular graph $G$ with $n$ vertices and $m$ edges in terms of $n$,$m$, $r$ and the signless Laplacian spectrum of $G$ with $x, y, z \in \{0, 1, +, -\}$.

## 2 Some preliminaries

Let $G$ be a graph with $V(G) = \{v_1, \ldots, v_n\}$ and $E(G) = \{e_1, \ldots, e_m\}$. The vertex-edge incidence matrix $R(G)$ of $G$ is the $n \times m$-matrix $(r_{ij})$, where $r_{ij} = 1$ if vertex $v_i$ is incident to edge $e_j$ and $r_{ij} = 0$ otherwise. Then $R(G)R(G)^\top = Q(G)$, $R(G)^\top R(G) = A(G^l) + 2I_m$. Let $q_1(G), q_2(G), \ldots, q_n(G)$ be the the signless Laplacian eigenvalues of $G$ arranged in non-decreasing order. Then $q_1 \geq 0$ and $R(G)^\top R(G)$ has eigenvalues 0 (of multiplicity $m-n$), $q_1, q_2, \ldots, q_n$.

For positive integers $p$ and $q$, let $J_{pq}$ be the all-ones $p \times q$-matrix, and in particular, let $J_p = J_{p,p}$.

In the rest of this paper, $G$ is an $r$-regular graph with $n$ vertices and $m$ edges. Then $2m = rn$. We write $A(G) = A$, $Q(G) = Q$, $R(G) = R$, and $q_i(G) = q_i$ for $i = 1, 2, \ldots, n$. In particular, $q_n = 2r$.

**Lemma 2.1** *Let $P(x, y)$ be a polynomial with two variables and real coefficients. Then*

*(1) $P(A, J_n)$ has the eigenvalues $P(r, n)$ and $P(q_i - r, 0)$ for $i = 1, 2, \ldots, n-1$, or equivalently, $P(Q, J_n)$ has the eigenvalues $P(2r, n)$ and $P(q_i, 0)$ for $i = 1, 2, \ldots, n-1$, and*

*(2) $P(R^\top R, J_m)$ has the eigenvalues $\sigma_m = P(2r, m)$ and $\sigma_i = P(q_i', 0)$ for $i = 1, 2, \ldots, m-1$, where $q_i' = 0$ for $i = 1, 2, \ldots, m-n$, and $q_i' = q_{i-m+n}$ for $i = m-n+1, \ldots, m-1$.*

*Proof.* (1) Let $X_1, X_2, \ldots, X_n$ be orthogonal eigenvectors of $A$ such that $AX_i = (q_i - r)X_i$ for $i = 1, 2, \ldots, n$. Since $A$ has equal row sums, $X_n = J_{n1}$. Since $J_n^2 = nJ_n$, $J_n J_{n1} = nJ_{n1}$ and $AJ_{n1} = rJ_{n1}$, we have $A^s J_n^t J_{n1} = A^s n^t J_{n1} = r^s n^t J_{n1}$ for nonnegative integers $s$ and $t$, and thus $P(Q, J_{nn})X_n = P(r, n)X_n$. For $i = 1, 2, \ldots, n-1$, since $J_n X_i = 0$ and $A^s X_i = (q_i - r)^s X_i$, we have $P(A, J_n)X_i = P(q_i - r, 0)X_i$.

(2) Since $R^\top R = A(G^l) + 2I_m$, $R(G)^\top R(G)$ have equal row sums. Note that $(R^\top R)^s J_m^t J_{m1} = (R^\top R)^s m^t J_{m1} = (2r)^s n^t J_{n1}$ for nonnegative integers $s$ and $t$. By similar argument as in (1), the result follows. □

**Lemma 2.2** *Let $B$ and $C$ be square matrices. Then*

$$\begin{vmatrix} B & E \\ F & C \end{vmatrix} = \begin{cases} |B||C - FB^{-1}E| & \textit{if } B \textit{ is invertible,} \\ |C||B - EC^{-1}F| & \textit{if } C \textit{ is invertible.} \end{cases}$$

Note that $q_{n-i}(G^c) = n - 2 - q_i$ for every $i \in \{1, 2, \ldots, n-1\}$ and $q_n(G^c) = 2(n - r - 1)$. Equivalently, we have the following lemma.

**Lemma 2.3** $(\lambda - n + 2 + 2r)f(x, G^c) = (-1)^n(\lambda - 2n + 2 + 2r)f(n - 2 - \lambda, G)$.

## 3 Signless Laplacian characteristic polynomials of $G^{xyz}$ with $z = 0$

Obviously, $f(\lambda, G^0) = \lambda^n$, $f(\lambda, G^+) = f(\lambda, G)$ and $f(\lambda, G^1) = (\lambda - 2n + 2)(\lambda - n + 2)^{n-1}$. Note that $G^l$ is regular of degree $2r - 2$ and $f(G^l, \lambda) = (\lambda - 2r + 4)^{m-n} f(\lambda - 2r + 4, G)$.

For $x, y \in \{0, 1, +, -\}$, $G^{xy0}$ consists of vertex disjoint $G^x$ and $(G^l)^y$), and then $f(\lambda, G^{xy0}) = f(\lambda, G^x) f(\lambda, (G^l)^y)$. Thus we have the following conclusions (16 cases).

$$\begin{aligned}
f(\lambda, G^{x00}) &= \lambda^m f(\lambda, G^x) \text{ for } x \in \{0, 1, +\},\\
f(\lambda, G^{x10}) &= (\lambda - 2m + 2)(\lambda - m + 2)^{m-1} f(\lambda, G^x) \text{ for } x \in \{0, 1, +\},\\
f(\lambda, G^{-00}) &= (-1)^n \lambda^m (\lambda - n + 2 + 2r)^{-1}\\
&\quad (\lambda - 2n + 2 + 2r) f(n - 2 - \lambda, G),\\
f(\lambda, G^{-10}) &= (-1)^n (\lambda - 2m + 2)(\lambda - m + 2)^{m-1} (\lambda - n + 2 + 2r)^{-1}\\
&\quad (\lambda - 2n + 2 + 2r) f(n - 2 - \lambda, G)\\
f(\lambda, G^{0+0}) &= \lambda^n (\lambda - 2r + 4)^{m-n} f(\lambda - 2r + 4, G^x),\\
f(\lambda, G^{1+0}) &= (\lambda - 2n + 2)(\lambda - n + 2)^{n-1} (\lambda - 2r + 4)^{m-n} f(\lambda - 2r + 4, G^x),\\
f(\lambda, G^{++0}) &= (\lambda - 2r + 4)^{m-n} f(\lambda, G) f(\lambda - 2r + 4, G^x),\\
f(\lambda, G^{-+0}) &= (-1)^n (\lambda - n + 2 + 2r)^{-1} (\lambda - 2n + 2 + 2r)\\
&\quad f(n - 2 - \lambda, G) f(\lambda - 2r + 4, G^x),\\
f(\lambda, G^{0-0}) &= (-1)^{n-1} \lambda^n (\lambda - m + 4r - 2)^{-1} (\lambda + 4r - 2m - 2)\\
&\quad (\lambda + 2r - 2 - m)^{m-n} f(m - 2r - \lambda + 2, G),\\
f(\lambda, G^{1-0}) &= (-1)^{n-1} (\lambda - 2n + 2)(\lambda - n + 2)^{n-1} (\lambda - m + 4r - 2)^{-1}\\
&\quad (\lambda + 4r - 2m - 2)(\lambda + 2r - 2 - m)^{m-n} f(m - 2r - \lambda + 2, G),\\
f(\lambda, G^{+-0}) &= (-1)^{n-1} (\lambda - m + 4r - 2)^{-1} (\lambda + 4r - 2m - 2)(\lambda + 2r - 2 - m)^{m-n}\\
&\quad f(\lambda, G) f(m - 2r - \lambda + 2, G),\\
f(\lambda, G^{--0}) &= -(\lambda - n + 2 + 2r)^{-1} (\lambda - 2n + 2 + 2r)(\lambda - m + 4r - 2)^{-1}\\
&\quad (\lambda + 4r - 2m - 2)(\lambda + 2r - 2 - m)^{m-n}\\
&\quad f(n - 2 - \lambda, G) f(m - 2r - \lambda + 2, G).
\end{aligned}$$

# 4 Signless Laplacian characteristic polynomials of $G^{xyz}$ with $z = 1$

Note that $G^{001}$ is a complete bipartite graph and $G^{111} = K_{n+m}$. Then

$$f(\lambda, G^{001}) = \lambda(\lambda - m - n)(\lambda - m)^{n-1}(\lambda - n)^{m-1},$$

$$f(\lambda, G^{111}) = (\lambda - 2n - 2m + 2)(\lambda - m - n + 2)^{m+n-1}.$$

**Theorem 4.1**

$$\begin{aligned} f(\lambda, G^{-01}) &= [(\lambda - n)(\lambda - 2n - m + 2r + 2) - mn](\lambda - n)^{m-1} \\ &\prod_{i=1}^{n-1}(\lambda - n - m + 2 + q_i), \end{aligned}$$

*Proof.* Obviously,

$$A(G^{-01}) = \begin{pmatrix} A(G^c) & J_{nm} \\ J_{mn} & 0 \end{pmatrix} = \begin{pmatrix} J_n - I_n - A & J_{nm} \\ J_{mn} & 0 \end{pmatrix}$$

and

$$D(G^{-01}) = \begin{pmatrix} (n + m - r - 1)I_n & 0 \\ 0 & nI_m \end{pmatrix}.$$

Then

$$\begin{aligned} f(\lambda, G^{-01}) &= \det(\lambda I_{n+m} - Q(G^{-01})) \\ &= \begin{vmatrix} (\lambda - n - m + r + 2)I_n - J_n + A & -J_{nm} \\ -J_{mn} & (\lambda - n)I_m \end{vmatrix}. \end{aligned}$$

Clearly, it is sufficient to prove our claim for $\lambda \neq n$. By Lemma 2.2,

$$f(\lambda, G^{-01}) = (\lambda - n)^{m-n}|B|.$$

where

$$B = (\lambda - n)(\lambda - n - m + r + 2)I_n - (\lambda - n)J_n + (\lambda - n)A - mJ_n.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\sigma_n = (\lambda - n)(\lambda - n - m + r + 2) - (\lambda - n)n + r(\lambda - n) - mn$$

$$= \quad (\lambda-n)(\lambda-2n-m+2r+2)-mn$$

and

$$\begin{aligned}\sigma_i &= (\lambda-n)(\lambda-n-m+r+2)+(\lambda-n)(q_i-r)\\ &= (\lambda-n)(\lambda-n-m+2+q_i)\end{aligned}$$

for $i=1,2,\ldots,n-1$. Then

$$|B| = [(\lambda-n)(\lambda-2n-m+2r+2)-mn](\lambda-n)^{n-1}\prod_{i=1}^{n-1}(\lambda-n-m+2+q_i),$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 4.2**

$$\begin{aligned}f(\lambda,G^{101}) &= [(\lambda-n)(\lambda-m-2n+2)-mn](\lambda-m-n+2)^{n-1}(\lambda-n)^{m-1},\\ f(\lambda,G^{+01}) &= [(\lambda-n)(\lambda-2r-m)-mn](\lambda-n)^{m-1}\prod_{i=1}^{n-1}(\lambda-m-q_i).\end{aligned}$$

**Theorem 4.3**

$$f(\lambda,G^{+11}) = [(\lambda-m-2r)(\lambda-2m-n+2)-mn](\lambda-m-n+2)^{m-1}\prod_{i=1}^{n-1}(\lambda-m-q_i).$$

*Proof.* Obviously,

$$A(G^{+11}) = \begin{pmatrix} A & J_{nm} \\ J_{mn} & J_m - I_m \end{pmatrix}$$

and

$$D(G^{+11}) = \begin{pmatrix} (m+r)I_n & 0 \\ 0 & (m+n-1)I_m \end{pmatrix}.$$

Then

$$f(\lambda,G^{+11}) \quad = \quad \det(\lambda I_{n+m} - Q(G^{+11}))$$

$$= \left| \begin{array}{cc} (\lambda-m-r)I_n - A & -J_{nm} \\ -J_{mn} & (\lambda-m-n+2)I_m - J_m \end{array} \right|.$$

Let

$$M = \begin{pmatrix} (\lambda-m-r)I_n - A & -J_{nm} \\ -J_{mn} & (\lambda-m-n+2)I_m - J_m \end{pmatrix}.$$

Obviously, $R^\top J_{nm} = 2J_m$. Thus multiplying the first row of the block matrix $M$ by $-\frac{1}{2}R^\top$ and adding the result to the second row of $M$, we obtain a new matrix

$$M_1 = \begin{pmatrix} (\lambda-m-r)I_n - A & -J_{nm} \\ \frac{m+r-\lambda}{2}R^\top + \frac{1}{2}R^\top A - J_{mn} & (\lambda-m-n+2)I_m \end{pmatrix}.$$

Clearly, $f(\lambda, G^{+11}) = |M| = |M_1|$ and it is sufficient to prove our claim for $\lambda \neq m+n-2$. By Lemma 2.2,

$$f(\lambda, G^{+11}) = (\lambda-m-n+2)^{m-n}|B|,$$

where

$$\begin{aligned} B &= (\lambda-m-r)(\lambda-n-m+2)I_n - (\lambda-m-n+2)A + \frac{m+r-\lambda}{2}J_{nm}R^\top \\ &\quad + \frac{1}{2}J_{nm}R^\top A - J_{nm}J_{mn}. \end{aligned}$$

Since $J_{nm}R^\top = rJ_n$, $J_{nm}R^\top A = r^2 J_n$ and $J_{nm}J_{mn} = mJ_n$, we have

$$B = (\lambda-m-r)(\lambda-n-m+2)I_n - (\lambda-m-n+2)A + \frac{(m+2r-\lambda)r}{2}J_n - mJ_n.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_n &= (\lambda-m-r)(\lambda-n-m+2) - (\lambda-m-n+2)r + \frac{(m+2r-\lambda)r}{2}n - mn \\ &= (\lambda-m-2r)(\lambda-n-2m+2) - mn \end{aligned}$$

and

$$\begin{aligned} \sigma_i &= (\lambda-m-r)(\lambda-n-m+2) - (\lambda-m-n+2)(q_i - r) \\ &= (\lambda-n-m+2)(\lambda-m-q_i). \end{aligned}$$

Then $|B| = [(\lambda-m-2r)(\lambda-n-2m+2) - mn](\lambda-n-m+2)^{n-1}\prod_{i=1}^{n-1}(\lambda-m-q_i)$, and thus the result follows. $\square$

Similarly, we have the following theorem.

**Theorem 4.4**

$$\begin{aligned} f(\lambda, G^{011}) &= [(\lambda - m)(\lambda - 2m - n + 2) - mn](\lambda - m)^{n-1}(\lambda - m - n + 2)^{m-1}, \\ f(\lambda, G^{-11}) &= [(\lambda - 2m - n + 2)(\lambda - m - 2n + 2r + 2) - mn](\lambda - m - n + 2)^{m-1} \\ &\quad \prod_{i=1}^{n-1}(\lambda - m - n + 2 + q_i). \end{aligned}$$

**Theorem 4.5**

$$\begin{aligned} f(\lambda, G^{0+1}) &= [(\lambda - m)(\lambda - n - 4r + 4) - mn](\lambda - n - 2r + 4)^{m-n}(\lambda - m)^{n-1} \\ &\quad \prod_{i=1}^{n-1}(\lambda - n - 2r + 4 - q_i). \end{aligned}$$

*Proof.* Obviously,

$$A(G^{0+1}) = \begin{pmatrix} 0 & J_{nm} \\ J_{mn} & A(G^l) \end{pmatrix} = \begin{pmatrix} 0 & J_{nm} \\ J_{mn} & R^\top R - 2I_m \end{pmatrix}$$

and

$$D(G^{0+1}) = \begin{pmatrix} mI_n & 0 \\ 0 & (n + 2r - 2)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{0+1}) = \begin{vmatrix} (\lambda - m)I_n & -J_{nm} \\ -J_{mn} & (\lambda - n - 2r + 4)I_m - R^\top R \end{vmatrix}.$$

Clearly, it is sufficient to prove our claim for $\lambda \neq m$. By Lemma 2.2 (and the fact that $J_{mn}J_{nm} = nJ_m$), $f(\lambda, G^{0+1}) = (\lambda - m)^{n-m}|B|$, where

$$B = (\lambda - m)(\lambda - n - 2r + 4)I_m - (\lambda - m)R^\top R - nJ_m.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_m &= (\lambda - m)(\lambda - n - 2r + 4) - 2r(\lambda - m) - nm \\ &= (\lambda - m)(\lambda - n - 4r + 4) - nm, \end{aligned}$$

for $1 \leq j \leq m - n$,

$$\sigma_j = (\lambda - m)(\lambda - n - 2r + 4),$$

and for $m-n+1 \leq j \leq m-1$

$$\begin{aligned} \sigma_i &= (\lambda-m)(\lambda-n-2r+4)-(\lambda-m)q'_j \\ &= (\lambda-m)(\lambda-n-2r+4)-(\lambda-m)q_{j-m+n} \\ &= (\lambda-m)(\lambda-n-2r+4-q_{j-m+n}). \end{aligned}$$

Then

$$\begin{aligned} |B| &= [(\lambda-m)(\lambda-n-4r+4)-mn](\lambda-n-2r+4)^{m-n}(\lambda-m)^{m-1} \\ & \prod_{i=1}^{n-1}(\lambda-n-2r+4-q_i), \end{aligned}$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 4.6**

$$\begin{aligned} f(\lambda, G^{0-1}) &= [(\lambda-m)(\lambda-2m-n+4r-2)-mn](\lambda-m-n+2r-2)^{m-n} \\ & (\lambda-m)^{n-1}\prod_{i=1}^{n-1}(\lambda-m-n-2+2r+q_i). \end{aligned}$$

**Theorem 4.7**

$$\begin{aligned} f(\lambda, G^{1+1}) &= [(\lambda-2n-m+2)(\lambda-n-4r+4)-mn](\lambda-n-2r+4)^{m-n} \\ & (\lambda-n-m+2)^{n-1}\prod_{i=1}^{n-1}(\lambda-n-2r+4-q_i). \end{aligned}$$

*Proof.* Obviously,

$$A(G^{1+1}) = \begin{pmatrix} J_n - I_n & J_{nm} \\ J_{mn} & R^\top R - 2I_m \end{pmatrix}$$

and

$$D(G^{1+1}) = \begin{pmatrix} (m+n-1)I_n & 0 \\ 0 & (n+2r-2)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{1+1}) \;=\; \begin{vmatrix} (\lambda-m-n+2)I_n - J_n & -J_{nm} \\ -J_{mn} & (\lambda-n-2r+4)I_m - R^{\top}R \end{vmatrix}.$$

Clearly, it is sufficient to prove our claim for $\lambda \neq m+2n-2$. Let

$$M = \begin{pmatrix} (\lambda-m-n+2)I_n - J_n & -J_{nm} \\ -J_{mn} & (\lambda-n-2r+4)I_m - R^{\top}R \end{pmatrix}.$$

Multiplying the first row of the block matrix $M$ by $\frac{1}{\lambda-2n-m+2}J_{mn}$ and adding the result to the second row of $M$, we obtain a new matrix

$$M_1 = \begin{pmatrix} (\lambda-m-n+2)I_n - J_n & -J_{nm} \\ 0 & \frac{-n}{\lambda-2n-m+2}J_m + (\lambda-n-2r+4)I_m - R^{\top}R \end{pmatrix},$$

Then

$$\begin{aligned} f(\lambda, G^{+11}) &= |M_1| \\ &= |(\lambda-m-n+2)I_n - J_n| \\ &\quad \cdot \left|(\lambda-n-2r+4)I_m - R^{\top}R + \frac{-n}{\lambda-2n-m+2}J_m\right| \\ &= (\lambda-n-m+2)^{n-1}(\lambda-2n-m+2)^{1-m}|B|, \end{aligned}$$

where

$$B = (\lambda-2n-m+2)(\lambda-n-2r+4)I_m - (\lambda-2n-m+2)R^{\top}R - nJ_m.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_m &= (\lambda-2n-m+2)(\lambda-n-2r+4) - (\lambda-2n-m+2)\cdot 2r - mn \\ &= (\lambda-2n-m+2)(\lambda-n-4r+4) - mn \end{aligned}$$

for $1 \leq j \leq m-n$,

$$\sigma_j = (\lambda-2n-m+2)(\lambda-n-2r+4)$$

and for $m-n+1 \leq j \leq m-1$,

$$\begin{aligned} \sigma_j &= (\lambda-2n-m+2)(\lambda-n-2r+4) - (\lambda-2n-m+2)q'_j \\ &= (\lambda-2n-m+2)(\lambda-n-2r+4) - (\lambda-2n-m+2)q_{j-m+n} \end{aligned}$$

$$= (\lambda - 2n - m + 2)(\lambda - n - 2r + 4 - q_{j-m+n}).$$

Then

$$\begin{aligned} |B| &= [(\lambda - 2n - m + 2)(\lambda - n - 4r + 4) - mn](\lambda - n - 2r + 4)^{m-n} \\ &\quad (\lambda - 2n - m + 2)^{m-1} \prod_{i=1}^{n-1} (\lambda - n - 2r + 4 - q_i), \end{aligned}$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 4.8**

$$\begin{aligned} f(\lambda, G^{++1}) &= [(\lambda - 2r - m)(\lambda - n - 4r + 4) - mn](\lambda - n - 2r + 4)^{m-n} \\ &\quad \prod_{i=1}^{n-1} (\lambda - n - 2r + 4 - q_i)(\lambda - m - q_i), \\ f(\lambda, G^{-+1}) &= [(\lambda - 2n - m + 2r + 2)(\lambda - n - 4r + 4) - mn] \\ &\quad (\lambda - n - 2r + 4)^{m-n} \\ &\quad \prod_{i=1}^{n-1} (\lambda - n - 2r + 4 - q_i)(\lambda - n - m + 2 + q_i), \\ f(\lambda, G^{1-1}) &= [(\lambda - 2n - m + 2)(\lambda - 2m - n + 4r - 2) - mn] \\ &\quad (\lambda - m - n - 2 + 2r)^{m-n} \\ &\quad (\lambda - n - m + 2)^{n-1} \prod_{i=1}^{n-1} (\lambda - m - n - 2 + 2r + q_i), \\ f(\lambda, G^{+-1}) &= [(\lambda - 2r - m)(\lambda - 2m - n + 4r - 2) - mn] \\ &\quad (\lambda - m - n - 2 + 2r)^{m-n} \\ &\quad \prod_{i=1}^{n-1} (\lambda - m - q_i)(\lambda - m - n - 2 + 2r + q_i), \\ f(\lambda, G^{--1}) & \\ &= [(\lambda - 2n - m + 2r + 2)(\lambda - 2m - n + 4r - 2) - mn] \\ &\quad (\lambda - m - n + 2r - 2)^{m-n} \\ &\quad \prod_{i=1}^{n-1} (\lambda - m - n + 2 + q_i)(\lambda - n - m + 2r - 2 + q_i). \end{aligned}$$

# 5 Signless Laplacian characteristic polynomials of $G^{xyz}$ with $z = +$

Since $G^{00+}$ is a bipartite graph, the signless Laplacian eigenvalues of $G^{00+}$ are the same as the Laplacian eigenvalues of $G^{00+}$. Thus

$$f(\lambda, G^{00+}) = \lambda(\lambda - r - 2)(\lambda - 2)^{m-n} \prod_{i=1}^{n-1} [(\lambda - 2)(\lambda - r) - q_i].$$

**Theorem 5.1**

$$f(\lambda, G^{10+}) = [\lambda^2 - (r + 2n)\lambda + 4n - 4](\lambda - 2)^{m-n} \prod_{i=1}^{n-1} [(\lambda - r - n + 2)(\lambda - 2) - q_i].$$

*Proof.* Obviously,

$$A(G^{10+}) = \begin{pmatrix} J_n - I_n & R \\ R^\top & 0 \end{pmatrix} \quad \textit{and} \quad D(G^{10+}) = \begin{pmatrix} (r + n - 1)I_n & 0 \\ 0 & 2I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{10+}) = \begin{vmatrix} (\lambda - r - n + 2)I_n - J_n & -R \\ -R^\top & (\lambda - 2)I_m \end{vmatrix}.$$

Clearly, it is sufficient to prove our claim for $\lambda \neq 2$. By Lemma 2.2 (and the fact $RR^\top = Q$),

$$\begin{aligned} f(\lambda, G^{10+}) &= |(\lambda - 2)I_m| \cdot \left| (\lambda - r - n + 2)I_n - J_n - \frac{1}{\lambda - 2} R I_m R^\top \right| \\ &= (\lambda - 2)^{m-n} \cdot |(\lambda - r - n + 2)(\lambda - 2)I_n - (\lambda - 2)J_n - Q|. \end{aligned}$$

Let $B = (\lambda - r - n + 2)(\lambda - 2)I_n - (\lambda - 2)J_n - Q$. By Lemma 2.1, the eigenvalues of $B$ are

$$\sigma_n = (\lambda - r - n + 2)(\lambda - 2) - (\lambda - 2)n - 2r = \lambda^2 - (r + 2n)\lambda + 4n - 4$$

and for $1 \leq i \leq n - 1$,

$$\sigma_i = (\lambda - r - n + 2)(\lambda - 2) - q_i.$$

Then

$$|B| = [\lambda^2 - (r + 2n)\lambda + 4n - 4] \prod_{i=1}^{n-1} [(\lambda - r - n + 2)(\lambda - 2) - q_i],$$

and thus the result follows. ☐

Similarly, we have the following theorem.

**Theorem 5.2**

$$\begin{aligned}
f(\lambda, G^{+0+}) &= [\lambda^2 - (2+3r)\lambda + 4r](\lambda-2)^{m-n}\prod_{i=1}^{n-1}[(\lambda-2)(\lambda-r-q_i)-q_i], \\
f(\lambda, G^{-0+}) &= [(\lambda-2)(\lambda-2n+r+2)-2r](\lambda-2)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda-2)(\lambda-n-r+2+q_i)-q_i].
\end{aligned}$$

**Theorem 5.3**

$$f(\lambda, G^{01+}) = [(\lambda-r)(\lambda-2m)-2r](\lambda-m)^{m-n}\prod_{i=1}^{n-1}[(\lambda-r)(\lambda-m)-q_i].$$

*Proof.* Obviously,

$$A(G^{01+}) = \begin{pmatrix} 0 & R \\ R^\top & J_m - I_m \end{pmatrix} \quad \textit{and} \quad D(G^{01+}) = \begin{pmatrix} rI_n & 0 \\ 0 & (m+1)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{01+}) = \begin{vmatrix} (\lambda-r)I_n & -R \\ -R^\top & (\lambda-m)I_m - J_m \end{vmatrix}.$$

Let

$$M = \begin{pmatrix} (\lambda-r)I_n & -R \\ -R^\top & (\lambda-m)I_m - J_m \end{pmatrix}.$$

Multiplying the first row of the block matrix $M$ by $-\frac{1}{2}J_{mn}$ and adding the result to the second row of $M$, we obtain a new matrix

$$M_1 = \begin{pmatrix} (\lambda-r)I_n & -R \\ \frac{r-\lambda}{2}J_{mn} - R^\top & (\lambda-m)I_m \end{pmatrix}.$$

Obviously, $f(\lambda, G^{01+}) = |M| = |M_1|$, and it is sufficient to prove our claim for $\lambda \neq m$. By Lemma 2.2,

$$f(\lambda, G^{01+}) = (\lambda-m)^{m-n}\left|(\lambda-r)(\lambda-m)I_n + \frac{r-\lambda}{2}RJ_{mn} - RR^\top\right|.$$

Let $B = (\lambda - r)(\lambda - m)I_n + \frac{r-\lambda}{2}RJ_{mn} - RR^{\top}$. Obviously, $RJ_{mn} = rJ_n$. Thus

$$B = (\lambda - r)(\lambda - m)I_n + \frac{r-\lambda}{2}rJ_n - Q.$$

By Lemma 2.1 (and the fact that $rn = 2m$), the eigenvalues of $B$ are

$$\sigma_n = (\lambda - r)(\lambda - m) + \frac{r-\lambda}{2}rn - 2r = (\lambda - r)(\lambda - 2m) - 2r$$

and for $1 \leq i \leq n-1$,

$$\sigma_i = (\lambda - r)(\lambda - m) - q_i.$$

Then

$$|B| = [(\lambda - r)(\lambda - 2m) - 2r]\prod_{i=1}^{n-1}[(\lambda - r)(\lambda - m) - q_i],$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 5.4**

$$\begin{aligned}
f(\lambda, G^{11+}) &= [(\lambda - r - 2n + 2)(\lambda - 2m) - 2r](\lambda - m)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - r - n + 2)(\lambda - m) - q_i], \\
f(\lambda, G^{-1+}) &= [(\lambda - 2m)(\lambda - 2n + r + 2) - 2r](\lambda - m)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - m)(\lambda - n + r + 2 - q_i) - q_i], \\
f(\lambda, G^{+1+}) &= [(\lambda - 2m)(\lambda - 3r) - 2r](\lambda - m)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - m)(\lambda - r - q_i) - q_i].
\end{aligned}$$

**Theorem 5.5**

$$\begin{aligned}
f(\lambda, G^{+++}) &= [(\lambda - 3r + 2)(\lambda - 4r)](\lambda - 2r + 2)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - r - q_i)(\lambda - 2r + 2 - q_i) - q_i].
\end{aligned}$$

*Proof.* Obviously,

$$A(G^{+++}) = \begin{pmatrix} A & R \\ R^\top & A(G^l) \end{pmatrix} \quad and \quad D(G^{+++}) = \begin{pmatrix} 2rI_n & 0 \\ 0 & 2rI_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{+++}) \;=\; \begin{vmatrix} (\lambda - 2r)I_n - A & -R \\ -R^\top & (\lambda - 2r + 2)I_m - R^\top R \end{vmatrix}.$$

Let

$$M = \begin{pmatrix} (\lambda - 2r)I_n - A & -R \\ -R^\top & (\lambda - 2r + 2)I_m - R^\top R \end{pmatrix}.$$

Multiplying the first row of the block matrix $M$ by $-R^\top$ and adding the result to the second row of $M$, we obtain a new matrix

$$M_1 \;=\; \begin{pmatrix} (\lambda - 2r)I_n - A & -R \\ (2r - \lambda - 1)R^\top + R^\top A & (\lambda - 2r + 2)I_m \end{pmatrix}.$$

Obviously, $f(\lambda, G^{++1}) = |M| = |M_1|$, and it is sufficient to prove our claim for $\lambda \neq 2r - 2$. By Lemma 2.2,

$$\begin{aligned} & f(\lambda, G^{+++}) \\ = \;& |(\lambda - 2r + 2)I_m| \cdot \left| (\lambda - 2r)I_n - A + \frac{1}{\lambda - 2r + 2} R((2r - \lambda - 1)R^\top + R^\top A) \right| \\ = \;& (\lambda - 2r + 2)^{m-n} \\ & |(\lambda - 2r)(\lambda - 2r + 2)I_n - (\lambda - 2r + 2)A + (2r - \lambda - 1)RR^\top + RR^\top A|. \end{aligned}$$

Let $B = (\lambda - 2r)(\lambda - 2r + 2)I_n - (\lambda - 2r + 2)A + (2r - \lambda - 1)RR^\top + RR^\top A$. Note that $RR^\top = Q$ and $A = Q - rI$. Then

$$B = (\lambda - 2r)(\lambda - 2r + 2)I_n - (\lambda - 2r + 2)(Q - rI) + (2r - \lambda - 1)Q + Q(Q - rI).$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\sigma_n = (\lambda - 2r)(\lambda - 2r + 2) - r(\lambda - 2r + 2) + 2r(2r - \lambda - 1) + 2r^2 = (\lambda - 3r + 2)(\lambda - 4r)$$

and for $1 \leq i \leq n - 1$,

$$\begin{aligned} \sigma_i &= (\lambda - 2r)(\lambda - 2r + 2) - (\lambda - 2r + 2)(q_i - r) + (2r - \lambda - 1)q_i + q_i(q_i - r) \\ &= (\lambda - r - q_i)(\lambda - 2r + 2 - q_i) - q_i. \end{aligned}$$

Then

$$|B| = [(\lambda - 3r + 2)(\lambda - 4r)] \prod_{i=1}^{n-1} [(\lambda - r - q_i)(\lambda - 2r + 2 - q_i) - q_i],$$

and thus the result follows. $\square$

Similarly, we can prove the following theorem.

**Theorem 5.6**

$$\begin{aligned} f(\lambda, G^{0++}) &= [(\lambda - r)(\lambda - 4r + 2) - 2r](\lambda - 2r + 2)^{m-n} \\ &\quad \prod_{i=1}^{n-1} [(\lambda - r)(\lambda - 2r + 2 - q_i) - q_i], \\ f(\lambda, G^{1++}) &= [(\lambda - r - 2n + 2)(\lambda - 4r + 2) - 2r](\lambda - 2r + 2)^{m-n} \\ &\quad \cdot \prod_{i=1}^{n-1} [(\lambda - r - n + 2)(\lambda - 2r + 2 - q_i) - q_i], \\ f(\lambda, G^{-++}) &= [(\lambda - 2n + r + 2)(\lambda - 4r + 2) - 2r](\lambda - 2r + 2)^{m-n} \\ &\quad \prod_{i=1}^{n-1} [(\lambda - n - r + 2 + q_i)(\lambda - 2r + 2 - q_i) - q_i]. \end{aligned}$$

**Theorem 5.7**

$$\begin{aligned} f(\lambda, G^{+-+}) &= [(\lambda - 3r)(\lambda - 2m + 4r - 4) - 2r](\lambda - m + 2r - 4)^{m-n} \\ &\quad \prod_{i=1}^{n-1} [(\lambda - r - q_i)(\lambda - m + 2r - 4 + q_i) - q_i]. \end{aligned}$$

*Proof.* Obviously,

$$A(G^{+-+}) = \begin{pmatrix} A & R \\ R^\top & J_m - I_m - A(G^l) \end{pmatrix} = \begin{pmatrix} A & R \\ R^\top & J_m + I_m - R^\top R \end{pmatrix}$$

and

$$D(G^{+-+}) = \begin{pmatrix} 2rI_n & 0 \\ 0 & (m - 2r + 3)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{+-+}) = \begin{vmatrix} (\lambda - 2r)I_n - A & -R \\ -R^\top & (\lambda - m + 2r - 4)I_m - J_m + R^\top R \end{vmatrix}.$$

Let

$$M=\begin{pmatrix} (\lambda-2r)I_n-A & -R \\ -R^\top & (\lambda-m+2r-4)I_m-J_m+R^\top R \end{pmatrix}.$$

Multiplying the first row of the block matrix $M$ by $-\frac{1}{2}J_{mn}+R^\top$ and adding the result to the second row of $M$, we obtain a new matrix

$$M_1 \;=\; \begin{pmatrix} (\lambda-2r)I_n-A & -R \\ (\lambda-2r-1)R^\top-R^\top A+\frac{3r-\lambda}{2}J_{mn} & (\lambda-m+2r-4)I_m \end{pmatrix}.$$

Obviously, $f(\lambda,G^{+-+})=|M|=|M_1|$, and it is sufficient to prove our claim for $\lambda\neq m-2r+4$. By Lemma 2.2,

$$f(\lambda,G^{+-+})=(\lambda-m+2r-4)^{m-n}\cdot|B|.$$

where $B=(\lambda-2r)(\lambda-m+2r-4)I_n-(\lambda-m+2r-4)A+(\lambda-2r-1)RR^\top+\frac{3r-\lambda}{2}RJ_{mn}-RR^\top A$. Note that $RR^\top=Q$, $RJ_{mn}=rJ_n$ and $A=Q-rI$. Then

$$\begin{aligned} B &= (\lambda-2r)(\lambda-m+2r-4)I_n-(\lambda-m+2r-4)(Q-rI)+(\lambda-2r-1)Q \\ &\quad +\frac{(3r-\lambda)r}{2}J_n-Q(Q-rI). \end{aligned}$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_n &= (\lambda-2r)(\lambda-m+2r-4)-r(\lambda-m+2r-4)+2r(\lambda-2r-1) \\ &\quad +\frac{(3r-\lambda)rn}{2}-2r^2 \\ &= (\lambda-3r)(\lambda-2m+4r-4)-2r \end{aligned}$$

and for $i=1,2,\ldots,n-1$

$$\begin{aligned} \sigma_i &= (\lambda-2r)(\lambda-m+2r-4)-(\lambda-m+2r-4)(q_i-r) \\ &\quad +(\lambda-2r-1)q_i-q_i(q_i-r) \\ &= (\lambda-r-q_i)(\lambda-m+2r-4+q_i)-q_i. \end{aligned}$$

Then

$$\begin{aligned} |B| &= [(\lambda-3r)(\lambda-2m+4r-4)-2r] \\ &\quad \prod_{i=1}^{n-1}[(\lambda-r-q_i)(\lambda-m+2r-4+q_i)-q_i], \end{aligned}$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 5.8**

$$
\begin{aligned}
f(\lambda, G^{0-+}) &= [(\lambda-r)(\lambda-2m+4r-4)-2r](\lambda-m+2r-4)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda-r)(\lambda-m+2r-4+q_i)-q_i], \\
f(\lambda, G^{1-+}) &= [(\lambda-r-2n+2)(\lambda-2m+4r-4)-2r](\lambda-m+2r-4)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda-r-n+2)(\lambda-m+2r-4+q_i)-q_i], \\
f(\lambda, G^{--+}) &= [(\lambda-2n+r+2)(\lambda-2m+4r-4)-2r](\lambda-m+2r-4)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda-n-r+2+q_i)(\lambda-m+2r-4+q_i)-q_i].
\end{aligned}
$$

# 6 Signless Laplacian characteristic polynomials of $G^{xyz}$ with $z=-$

Since $G^{00-}$ is a bipartite graph, the signless laplacian eigenvalues of $G^{00-}$ are the same as the laplacian eigenvalues of it.

$$
f(\lambda, G^{00-}) = \lambda(\lambda-n-m+r+2)(\lambda-n+2)^{m-n}\prod_{i=1}^{n-1}\{(\lambda-m+r)(\lambda-n+2)-q_i\}.
$$

**Theorem 6.1**

$$
\begin{aligned}
f(\lambda, G^{10-}) &= [(\lambda-n+2)(\lambda-2n+m+r+2)+(2r-m)n-2r] \\
&\quad (\lambda-n+2)^{m-n}\prod_{i=1}^{n-1}[(\lambda-m-n+r+2)(\lambda-n+2)-q_i].
\end{aligned}
$$

*Proof.* Obviously,

$$
A(G^{10-}) = \begin{pmatrix} J_n - I_n & J_{nm} - R \\ J_{mn} - R^\top & 0 \end{pmatrix}
$$

and

$$
D(G^{10-}) = \begin{pmatrix} (n+m-r-1)I_n & 0 \\ 0 & (n-2)I_m \end{pmatrix}.
$$

Then

$$f(\lambda, G^{10-}) = \begin{vmatrix} (\lambda - n - m + r + 2)I_n - J_n & R - J_{nm} \\ R^\top - J_{mn} & (\lambda - n + 2)I_m \end{vmatrix}.$$

Clearly, it is sufficient to prove our claim for $\lambda \neq n - 2$. By Lemma 2.2,

$$f(\lambda, G^{10-}) = (\lambda - n + 2)^{m-n}|B|,$$

where

$$\begin{aligned} B &= (\lambda - n - m + r + 2)(\lambda - n + 2)I_n - (\lambda - n + 2)J_n \\ &\quad -(R - J_{nm})(R^\top - J_{mn}). \end{aligned}$$

Note that $RR^\top = Q$, $J_{nm}R^\top = rJ_n$ and $RJ_{mn} = rJ_n$, then

$$B = (\lambda - n - m + r + 2)(\lambda - n + 2)I_n + (2r + n - m - 2 - \lambda)J_n - Q.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_n &= (\lambda - n - m + r + 2)(\lambda - n + 2) + (2r + n - m - 2 - \lambda)n - 2r \\ &= (\lambda - n + 2)(\lambda - 2n + m + r + 2) + (2r - m)n - 2r \end{aligned}$$

and for $i = 1, 2, \ldots, n - 1$,

$$\sigma_i = (\lambda - n - m + r + 2)(\lambda - n + 2) - q_i.$$

Then

$$\begin{aligned} |B| &= [(\lambda - n + 2)(\lambda - 2n + m + r + 2) + (2r - m)n - 2r] \\ &\quad \prod_{i=1}^{n-1}[(\lambda - m - n + r + 2)(\lambda - n + 2) - q_i], \end{aligned}$$

and thus the result follows. $\square$

Similarly, we have the following theorem.

**Theorem 6.2**

$$\begin{aligned} f(\lambda, G^{+0-}) &= [(\lambda - n + 2)(\lambda - m - r) + (2r - m)n - 2r](\lambda - n + 2)^{m-n} \\ &\quad \prod_{i=1}^{n-1}[(\lambda - n + 2)(\lambda - m + r - q_i) - q_i], \\ f(\lambda, G^{-0-}) &= [(\lambda - n + 2)(\lambda - 2n - m + 3r + 2) + (2r - m)n - 2r](\lambda - n + 2)^{m-n} \\ &\quad \prod_{i=1}^{n-1}[(\lambda - n + 2)(\lambda - n - m + r + 2 + q_i) - q_i]. \end{aligned}$$

**Theorem 6.3**

$$\begin{aligned} f(\lambda, G^{01-}) &= [(\lambda - m + r)(\lambda - n - 2m + 4) + (4 - n)m - 2r](\lambda - n - m + 4)^{m-n} \\ &\quad \prod_{i=1}^{n-1}[(\lambda - m - n + 4)(\lambda - m + r) - q_i]. \end{aligned}$$

*Proof.* Obviously,

$$A(G^{01-}) = \begin{pmatrix} 0 & J_{nm} - R \\ J_{mn} - R^{\top} & J_m - I_m \end{pmatrix}$$

and

$$D(G^{01-}) = \begin{pmatrix} (m - r)I_n & 0 \\ 0 & (m + n - 3)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{01-}) = \begin{vmatrix} (\lambda - m + r)I_n & R - J_{nm} \\ R^{\top} - J_{mn} & (\lambda - n - m + 4)I_m - J_m \end{vmatrix}.$$

Clearly, it is sufficient to prove our claim for $\lambda \neq m - r$. By Lemma 2.2 and the fact $R^{\top} J_{nm} = 2J_n$ and $J_{mn}R = 2J_n$,

$$\begin{aligned} & f(\lambda, G^{01-}) \\ = \; & (\lambda - m + r)^{n-m} \\ & |(\lambda - n - m + 4)(\lambda - m + r)I_m - (\lambda - m + r)J_m - (R^{\top} - J_{mn})(R - J_{nm})| \\ = \; & (\lambda - m + r)^{n-m} \\ & |(\lambda - n - m + 4)(\lambda - m + r)I_m - R^{\top}R + (m + 4 - \lambda - r - n)J_m| \end{aligned}$$

Let $B = (\lambda - n - m + 4)(\lambda - m + r)I_m - R^{\top}R + (m + 4 - \lambda - r - n)J_m$. By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_n &= (\lambda - n - m + 4)(\lambda - m + r) - 2r + (m + 4 - \lambda - r - n)m \\ &= (\lambda - m + r)(\lambda - n - 2m + 4) + (4 - n)m - 2r, \end{aligned}$$

for $1 \leq j \leq m - n$,

$$\sigma_i = (\lambda - n - m + 4)(\lambda - m + r),$$

and for $m - n + 1 \leq j \leq m - 1$,

$$\sigma_j = (\lambda - n - m + 4)(\lambda - m + r) - q'_j = (\lambda - n - m + 4)(\lambda - m + r) - q_{j-m+n}.$$

Then

$$\begin{aligned}|B| &= [(\lambda-m+r)(\lambda-n-2m+4)+(4-n)m-2r](\lambda-m+r)^{m-n}\\ &\quad (\lambda-n-m+4)^{m-n}\prod_{i=1}^{n-1}[(\lambda-m-n+4)(\lambda-m+r)-q_i].\end{aligned}$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 6.4**

$$\begin{aligned}f(\lambda,G^{0+-}) &= [(\lambda-m+r)(\lambda-n-4r+6)+(4-n)m-2r](\lambda-n-2r+6)^{m-n}\\ &\quad \prod_{i=1}^{n-1}[(\lambda-m+r)(\lambda-n-2r+6-q_i)-q_i],\\ f(\lambda,G^{0--}) &= [(\lambda-m+r)(\lambda-n-2m+4r)+(4-n)m-2r](\lambda-n-m+2r)^{m-n}\\ &\quad \prod_{i=1}^{n-1}[(\lambda-m+r)(\lambda-n-m+2r+q_i)-q_i].\end{aligned}$$

**Theorem 6.5**

$$\begin{aligned}f(\lambda,G^{11-}) &= [(\lambda-2n-2m+2)(\lambda-n-m+r+4)+8m](\lambda-n-m+4)^{m-n}\\ &\quad \prod_{i=1}^{n-1}[(\lambda-m-n+r+2)(\lambda-n-m+4)-q_i].\end{aligned}$$

*Proof.* Obviously,

$$A(G^{11-}) = \begin{pmatrix} J_n-I_n & J_{nm}-R\\ J_{mn}-R^\top & J_m-I_m\end{pmatrix}$$

and

$$D(G^{11-}) = \begin{pmatrix} (n+m-r-1)I_n & 0\\ 0 & (m+n-3)I_m\end{pmatrix}.$$

Then

$$\begin{aligned}f(\lambda,G^{11-}) &= \begin{vmatrix} (\lambda-m-n+r+2)I_n-J_n & R-J_{nm}\\ R^\top-J_{mn} & (\lambda-n-m+4)I_m-J_m\end{vmatrix}\\ &= (2-n)^{-m}|M|,\end{aligned}$$

where

$$M = \begin{pmatrix} (\lambda - m - n + r + 2)I_n - J_n & R - J_{nm} \\ (2-n)R^\top - (2-n)J_{mn} & (2-n)(\lambda - n - m + 4)I_m - (2-n)J_m \end{pmatrix}$$

Obviously, $J_{mn}R = 2J_{mm}$ and $J_{mn}J_{nm} = nJ_{mm}$. Hence multiplying the first row of the block matrix $M$ by $J_{mn}$ and adding the result to the second row of $M$, we obtain a new matrix

$$M' = \begin{pmatrix} (\lambda - m - n + r + 2)I_n - J_n & R - J_{nm} \\ (2-n)R^\top + (\lambda - m - n + r)J_{mn} & (2-n)(\lambda - n - m + 4)I_m \end{pmatrix}.$$

Clearly, $|M'| = |M|$ and $f(\lambda, G^{11-}) = (2-n)^{-m}|M'|$. Obviously, it is sufficient to prove our claim for $\lambda \neq n + m - 4$. By Lemma 2.2, $f(\lambda, G^{11-}) = (2-n)^{-n}(\lambda - n - m + 4)^{m-n}|B|$, where

$$\begin{aligned} B &= (2-n)(\lambda - m - n + r + 2)(\lambda - n - m + 4)I_n \\ &\quad -(2-n)Q - (2-n)(\lambda - n - m + 4)J_n \\ &\quad -(\lambda - m + r - 2)rJ_n + (\lambda - n - m + r)mJ_n. \end{aligned}$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_n &= (2-n)(\lambda - m - n + r + 2)(\lambda - n - m + 4) - 2r(2-n) \\ &\quad -(2-n)(\lambda - n - m + 4)n \\ &\quad -(\lambda - m + r - 2)rn + (\lambda - n - m + r)mn \\ &= (2-n)[(\lambda - 2n - 2m + 2)(\lambda - n - m + 4 + r) + 8m], \end{aligned}$$

and for $i = 1, 2, \ldots, n-1$

$$\begin{aligned} \sigma_i &= (2-n)(\lambda - m - n + r + 2)(\lambda - n - m + 4) - (2-n)q_i \\ &= (2-n)[(\lambda - m - n + r + 2)(\lambda - n - m + 4) - q_i]. \end{aligned}$$

Then

$$\begin{aligned} |B| &= (2-n)^n[(\lambda - 2n - 2m + 2)(\lambda - n - m + r + 4) + 8m] \\ &\quad \prod_{i=1}^{n-1}[(\lambda - m - n + r + 2)(\lambda - n - m + 4) - q_i], \end{aligned}$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 6.6**

$$\begin{aligned} f(\lambda, G^{+1-}) &= [(\lambda-m-r)(\lambda-n-2m+4)+(4-n)m-2r] \\ &\quad (\lambda-n-m+4)^{m-n}\prod_{i=1}^{n-1}[(\lambda-n-m+4)(\lambda-m+r-q_i)-q_i], \\ f(\lambda, G^{-1-}) &= [(\lambda-n-2m+4)(\lambda-2n-m+3r+2)+(4-n)m-2r] \\ &\quad (\lambda-n-m+4)^{m-n}\prod_{i=1}^{n-1}[(\lambda-n-m+4)(\lambda-m-n+r+q_i+2)-q_i]. \end{aligned}$$

**Theorem 6.7**

$$\begin{aligned} f(\lambda, G^{1+-}) &= [(\lambda-2n-m+r+2)(\lambda-n-4r+6)+(4-n)m-2r] \\ &\quad (\lambda-n-2r+6)^{m-n}\prod_{i=1}^{n-1}[(\lambda-n-m+r+2)(\lambda-n-2r+6-q_i)-q_i]. \end{aligned}$$

*Proof.* Obviously,

$$A(G^{1+-}) = \begin{pmatrix} J_n - I_n & J_{nm} - R \\ J_{mn} - R^\top & A(G^l) \end{pmatrix} = \begin{pmatrix} J_n - I_n & J_{nm} - R \\ J_{mn} - R^\top & R^\top R - 2I_m \end{pmatrix}$$

and

$$D(G^{1+-}) = \begin{pmatrix} (n+m-r-1)I_n & 0 \\ 0 & (n+2r-4)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{1+-}) = \begin{vmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ R^\top - J_{mn} & (\lambda-n-2r+6)I_m - R^\top R \end{vmatrix}.$$

Let

$$M = \begin{pmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ R^\top - J_{mn} & (\lambda-n-2r+6)I_m - R^\top R \end{pmatrix},$$

then $f(\lambda, G^{1+-}) = |M|$. Obviously, $R^\top J_{nm} = 2J_m$. Thus multiplying the first row of the block matrix $M$ by $R^\top$ and adding the result to the second row of $M$, we obtain a new matrix

$$M' = \begin{pmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ (\lambda-n-m+r+3)R^\top - 3J_{mn} & (\lambda-n-2r+6)I_m - 2J_m \end{pmatrix}.$$

Let

$$M'' = \begin{pmatrix} (\lambda - n - m + r + 2)I_n - J_n & R - J_{nm} \\ (2-n)(\lambda - n - m + r + 3)R^\top & \\ -3(2-n)J_{mn} & (2-n)(\lambda - n - 2r + 6)I_m - 2(2-n)J_m \end{pmatrix}.$$

Clearly, $|M''| = (2-n)^m|M'| = (2-n)^m|M|$ and $f(\lambda, G^{11-}) = |M'| = (2-n)^{-m}|M''|$. Obviously, $J_{mn}R = 2J_m$ and $J_{mn}J_{nm} = nJ_m$. Thus multiplying the first row of the block matrix $M''$ by $2J_{mn}$ and adding the result to the second row of $M''$, we obtain a new matrix

$$M''' = \begin{pmatrix} (\lambda - n - m + r + 2)I_n - J_n & R - J_{nm} \\ (2-n)(\lambda - n - m + r + 3)R^\top & \\ +(2\lambda - n - 2m + 2r - 2)J_{mn} & (2-n)(\lambda - n - 2r + 6)I_m \end{pmatrix}.$$

Clearly, $|M'''| = |M''|$, $f(\lambda, G^{11-}) = (2-n)^{-m}|M'''|$, and it is sufficient to prove our claim for $\lambda \neq n + 2r - 6$. By Lemma 2.2,

$$|M'''| = (2-n)^{m-n}(\lambda - n - 2r + 6)^{m-n}|B|,$$

where

$$\begin{aligned} B &= (2-n)(\lambda - n - 2r + 6)(\lambda - n - m + r + 2)I_n - (2-n)(\lambda - n - 2r + 6)J_n \\ &\quad -(R - J_{nm})[(2-n)(\lambda - n - m + r + 3)R^\top + (2\lambda - n - 2m + 2r - 2)J_{mn}] \\ &= (2-n)(\lambda - n - 2r + 6)(\lambda - n - m + r + 2)I_n \\ &\quad -(2-n)(\lambda - n - m + r + 3)RR^\top \\ &\quad -(2-n)(\lambda - n - 2r + 6)J_n - (4n - 8 + \lambda n - n^2 - nm + nr)rJ_n \\ &\quad +(2\lambda - n - 2m + 2r - 2)mJ_n. \end{aligned}$$

Thus

$$f(\lambda, G^{11-}) = (2-n)^{-n}(\lambda - n - 2r + 6)^{m-n}|B|.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned} \sigma_n &= (2-n)(\lambda - n - 2r + 6)(\lambda - n - m + r + 2) - (2-n)(\lambda - n - m + r + 3) \cdot 2r \\ &\quad -(2-n)(\lambda - n - 2r + 6)n - (4n - 8 + \lambda n - n^2 - nm + nr)rn \\ &\quad + (2\lambda - n - 2m + 2r - 2)mn \\ &= (2-n)[(\lambda - 2n - m + r + 2)(\lambda - n - 4r + 6) + (4-n)m - 2r], \end{aligned}$$

and for $i = 1, 2, \ldots, n-1$,

$$\begin{aligned}
\sigma_i &= (2-n)(\lambda - n - 2r + 6)(\lambda - n - m + r + 2) - (2-n)(\lambda - n - m + r + 3)q_i \\
&= (2-n)[(\lambda - n - m + r + 2)(\lambda - n - 2r + 6 - q_i) - q_i].
\end{aligned}$$

Then

$$\begin{aligned}
|B| &= (2-n)^n[(\lambda - 2n - m + r + 2)(\lambda - n - 4r + 6) + (4-n)m - 2r] \\
&\quad \prod_{i=1}^{n-1}[(\lambda - n - m + r + 2)(\lambda - n - 2r + 6 - q_i) - q_i],
\end{aligned}$$

and thus the result follows. $\square$

Similarly, we have the following theorem.

**Theorem 6.8**

$$\begin{aligned}
f(\lambda, G^{++-}) &= [(\lambda - m - r)(\lambda - n - 4r + 6) + (4-n)m - 2r](\lambda - n - 2r + 6)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - m + r - q_i)(\lambda - n - 2r + 6 - q_i) - q_i], \\
f(\lambda, G^{-+-}) &= [(\lambda - n - 4r + 6)(\lambda - 2n - m + 3r + 2) + (4-n)m - 2r] \\
&\quad (\lambda - n - 2r + 6)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - m - n + r + 2 + q_i)(\lambda - n - 2r + 6 - q_i) - q_i].
\end{aligned}$$

**Theorem 6.9**

$$\begin{aligned}
f(\lambda, G^{1--}) &= [(\lambda - n - 2m + 4r)(\lambda - 2n - m + r + 2) + (4-n)m - 2r] \\
&\quad (\lambda - n - m + 2r)^{m-n} \\
&\quad \prod_{i=1}^{n-1}[(\lambda - n - m + r + 2)(\lambda - n - m + 2r + q_i) - q_i].
\end{aligned}$$

*Proof.* Obviously,

$$A(G^{1--}) = \begin{pmatrix} J_n - I_n & J_{nm} - R \\ J_{mn} - R^\top & J_m - I_m - A(G^l) \end{pmatrix} = \begin{pmatrix} J_n - I_n & J_{nm} - R \\ J_{mn} - R^\top & J_m + I_m - R^\top R \end{pmatrix}$$

and

$$D(G^{1--}) = \begin{pmatrix} (n+m-r-1)I_n & 0 \\ 0 & (n+m-2r-1)I_m \end{pmatrix}.$$

Then

$$f(\lambda, G^{1--}) \;=\; \begin{vmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ R^\top - J_{mn} & (\lambda-n-m+2r)I_m - J_m + R^\top R \end{vmatrix}.$$

Let

$$M = \begin{pmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ R^\top - J_{mn} & (\lambda-n-m+2r)I_m - J_m + R^\top R \end{pmatrix}.$$

Then $f(\lambda, G^{1--}) = |M|$. Note that $R^\top J_{nm} = 2J_m$ . Hence multiplying the first row of the block matrix $M$ by $-R^\top$ and adding the result to the second row of $M$, we obtain a new matrix

$$M_1 = \begin{pmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ (n+m-\lambda-r-1)R^\top + J_{mn} & (\lambda-n-m+2r)I_m + J_m \end{pmatrix}.$$

Let

$$M_2 = \begin{pmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ (2-n)(n+m-\lambda-r-1)R^\top & \\ +(2-n)J_{mn} & (2-n)(\lambda-n-m+2r)I_m + (2-n)J_m \end{pmatrix}.$$

Clearly, $|M_2| = (2-n)^{-m}|M_1| = (2-n)^{-m}|M|$ and $f(\lambda, G^{1--}) = |M_1| = (2-n)^{-m}|M_2|$. Obviously, $J_{mn}R = 2J_m$ and $J_{mn}J_{nm} = nJ_m$. Thus multiplying the first row of the block matrix $M''$ by $-J_{mn}$ and adding the result to the second row of $M_2$, we obtain a new matrix

$$M_3 = \begin{pmatrix} (\lambda-n-m+r+2)I_n - J_n & R - J_{nm} \\ (2-n)(n+m-\lambda-r-1)R^\top + (n+m-\lambda-r)J_{mn} & (2-n)(\lambda-n-m+2r)I_m \end{pmatrix}.$$

Clearly, $|M_3| = |M_2|$, $f(\lambda, G^{1--}) = (2-n)^{-m}|M_3|$, and it is sufficient to prove our claim for $\lambda \neq n+m-2r$. By Lemma 2.2,

$$|M_3| = (2-n)^{m-n}(\lambda-n-m+2r)^{m-n}|B|,$$

where

$$B \;=\; (2-n)(\lambda-n-m+2r)(\lambda-n-m+r+2)I_n - (2-n)(\lambda-n-m+2r)J_n$$

$$\begin{aligned}
&-(R-J_{nm})((2-n)(n+m-\lambda-r-1)R^\top+(n+m-\lambda-r)J_{mn})\\
=\;&(2-n)(\lambda-n-m+2r)(\lambda-n-m+r+2)I_n-(2-n)(n+m-\lambda-r-1)Q\\
&-(2-n)(\lambda-n-m+2r)J_n+(m+n\lambda+nr+2n-\lambda-r-n^2-nm-2)rJ_n\\
&+(n+m-\lambda-r)mJ_n.
\end{aligned}$$

Thus

$$f(\lambda,G^{1--})=(2-n)^{-n}(\lambda-n-m+2r)^{m-n}|B|.$$

By Lemma 2.1, the eigenvalues of $B$ are

$$\begin{aligned}
\sigma_n\;=\;&(2-n)(\lambda-n-m+2r)(\lambda-n-m+r+2)-(2-n)(n+m-\lambda-r-1)2r\\
&-(2-n)(\lambda-n-m+2r)n+(m+n\lambda+nr+2n-\lambda-r-n^2-nm-2)rn\\
&+(n+m-\lambda-r)mn\\
=\;&(2-n)[(\lambda-n-2m+4r)(\lambda-2n-m+r+2)+(4-n)m-2r],
\end{aligned}$$

and

$$\begin{aligned}
\sigma_i\;=\;&(2-n)(\lambda-n-m+2r)(\lambda-n-m+r+2)-(2-n)(n+m-\lambda-r-1)q_i\\
=\;&(2-n)[(\lambda-n-m+r+2)(\lambda-n-m+2r+q_i)-q_i]
\end{aligned}$$

for $i=1,2,\ldots,n-1$. Then

$$\begin{aligned}
|B|\;=\;&(2-n)^n[(\lambda-n-2m+4r)(\lambda-2n-m+r+2)+(4-n)m-2r]\\
&\prod_{i=1}^{n-1}[(\lambda-n-m+r+2)(\lambda-n-m+2r+q_i)-q_i],
\end{aligned}$$

and thus the result follows. □

Similarly, we have the following theorem.

**Theorem 6.10**

$$\begin{aligned}
f(\lambda,G^{+--})\;=\;&[(\lambda-m-r)(\lambda-n-2m+4r)+(4-n)m-2r]\\
&(\lambda-n-m+2r)^{m-n}\\
&\prod_{i=1}^{n-1}[(\lambda-n-m+2r+q_i)(\lambda+r-m-q_i)-q_i].
\end{aligned}$$

**Theorem 6.11**

$$\begin{aligned} f(\lambda, G^{---}) &= (\lambda-2n-2m+4r+2)(\lambda+3r-n-m)(\lambda+2r-n-m)^{m-n} \\ &\quad \prod_{i=1}^{n-1}[(\lambda-n-m+r+q_i+2)(\lambda+2r-n-m+q_i)-q_i]. \end{aligned}$$

*Proof.* Note that $G^{---}$ is the complement of $G^{+++}$, and $G^{+++}$ is $2r$-regular. By Lemma 2.3 and Theorem 5.5,

$$\begin{aligned} & f(\lambda, G^{---}) \\ = \;& (-1)^{m+n-1}\frac{\lambda-2m-2n+2+4r}{n+m-2-\lambda-4r}(n+m-2-\lambda-4r)(n+m-\lambda-3r) \\ & (m+n-\lambda-2r)^{m-n}\prod_{i=1}^{n-1}[(n+m-\lambda-r-q_i-2)(n+m-\lambda-2r-q_i)-q_i] \\ = \;& (\lambda-2n-2m+4r+2)(\lambda+3r-n-m)(\lambda+2r-n-m)^{m-n} \\ & \prod_{i=1}^{n-1}[(\lambda-n-m+r+q_i+2)(\lambda+2r-n-m+q_i)-q_i], \end{aligned}$$

as desired. □

**Acknowledgement.** This work was supported by the Research Fund for the Doctoral Program of Higher Education of China (No. 20124407110002).

# References

[1] D.M. Cvetković, M. Doob, H. Sachs, Spectra of Graphs: Theory and Applications, third ed., Johann Ambrosius Barth Verlag, Heidelberg, Leipzig, 1995.

[2] A.K. Kelmans, The properties of the characteristic polynomial of a graph, in: Cyberneticsin the Service of Communism 4 (Russian), Izdat. “Ѐnergija”, Moscow, 1967, pp. 27–41.

[3] J. Yan, K. Xu, Spectra of transformation graphs of regular graph, Appl. Math. J. Chinese Univ. Ser. A 23 (2008) 476-480.

[4] A. Deng, A. Kelmans, J. Meng, Laplacian spectra of regular graph transformation, Discrete Appl. Math. 161 (2013) 118–133.